

## Press Release

# ENHERTU<sup>®</sup> Granted Priority Review in the U.S. for Patients with Metastatic HER2 Positive Solid Tumors

- Application based on results from DESTINY-PanTumor02 trial and supported by additional ENHERTU data
- Submission to be reviewed under FDA Real Time Oncology Review and Project Orbis
- If approved, Daiichi Sankyo and AstraZeneca's ENHERTU will potentially be the first HER2 directed treatment and antibody drug conjugate to receive a tumor agnostic indication

**Tokyo and Basking Ridge, NJ – (January 29, 2024)** – Daiichi Sankyo (TSE: 4568) and AstraZeneca's (LSE/STO/Nasdaq: AZN) supplemental Biologics License Application (sBLA) for ENHERTU<sup>®</sup> (fam-trastuzumab deruxtecan-nxki) has been accepted and granted Priority Review in the U.S. for the treatment of adult patients with unresectable or metastatic HER2 positive (immunohistochemistry [IHC] 3+) solid tumors who have received prior treatment or who have no satisfactory alternative treatment options.

ENHERTU is a specifically engineered HER2 directed antibody drug conjugate (ADC) being jointly developed and commercialized by Daiichi Sankyo and AstraZeneca.

The U.S. Food and Drug Administration (FDA) grants Priority Review to applications for medicines that, if approved, would offer significant improvements over available treatment options by demonstrating safety or efficacy improvements, preventing serious conditions or enhancing patient compliance. The Prescription Drug User Fee Act (PDUFA) date, the FDA action date for their regulatory decision, is May 30, 2024. The Priority Review follows receipt of [Breakthrough Therapy Designation](#) granted by the FDA in August 2023 for ENHERTU in metastatic HER2 positive solid tumors.

The sBLA is being reviewed under the Real-Time Oncology Review (RTOR) program and Project Orbis, two initiatives of the FDA which are designed to bring safe and effective cancer treatments to patients as early as possible. RTOR allows the FDA to review components of an application before submission of the complete application. Project Orbis provides a framework for concurrent submission and review of oncology medicines among participating international partners.

The sBLA is based on data from the ongoing [DESTINY-PanTumor02](#) phase 2 trial where ENHERTU demonstrated clinically meaningful and durable responses leading to a clinically meaningful survival benefit in previously treated patients across HER2 expressing advanced solid tumors. Data from other supporting

trials in patients with HER2 positive IHC 3+ tumors in the ENHERTU clinical development program, including [DESTINY-Lung01](#) and [DESTINY-CRC02](#), also were included in the submission.

“The clinical benefit seen across HER2 expressing metastatic solid tumors in the DESTINY-PanTumor02 trial and ongoing data from the ENHERTU clinical development program continues to demonstrate the potential of this medicine beyond its approved indications,” said Ken Takeshita, MD, Global Head, R&D, Daiichi Sankyo. “If approved, ENHERTU could become the first HER2 directed therapy and antibody drug conjugate with a tumor agnostic indication, providing patients with a potential new treatment option.”

“Today’s Priority Review for the first tumor agnostic submission for ENHERTU reflects the potential of this medicine to redefine the treatment of HER2 expressing cancers,” said Susan Galbraith, MBBChir, PhD, Executive Vice President, Oncology R&D, AstraZeneca. “Biomarkers for HER2 expression are already established in breast and gastric cancers, but we must now define them across tumor types. We will continue working closely with the FDA to bring this potential first tumor agnostic HER2 targeted medicine to patients as quickly as possible.”

#### **About DESTINY-PanTumor02**

DESTINY-PanTumor02 is a global, multicenter, multi-cohort, open-label phase 2 trial evaluating the efficacy and safety of ENHERTU (5.4 mg/kg) for the treatment of previously treated HER2 expressing tumors, including biliary tract, bladder, cervical, endometrial, ovarian, pancreatic cancer or other tumors. The primary efficacy endpoint of DESTINY-PanTumor02 is confirmed objective response rate (ORR) as assessed by investigator. Secondary endpoints include duration of response (DOR), disease control rate (DCR), progression-free survival (PFS), overall survival (OS), safety, tolerability and pharmacokinetics. DESTINY-PanTumor02 enrolled 267 patients at multiple sites in Asia, Europe and North America. For more information about the trial, visit [ClinicalTrials.gov](#).

The primary analysis of this study was [presented](#) in a late-breaking mini-oral session at the 2023 European Society for Medical Oncology (ESMO) Congress and simultaneously published in the [Journal of Clinical Oncology](#).

#### **About DESTINY-Lung01**

DESTINY-Lung01 is a global phase 2, open-label, two-cohort trial evaluating the efficacy and safety of ENHERTU (6.4 mg/kg and 5.4 mg/kg) in patients with *HER2* mutant (cohort 2, n=91) or *HER2* overexpressing (cohort 1 and 1a, n=90) (defined as IHC 3+ or IHC 2+) unresectable or metastatic non-squamous non-small cell lung cancer (NSCLC) who had progressed after one or more systemic therapies. The primary endpoint is confirmed ORR by independent central review. Key secondary endpoints include

DOR, DCR, PFS, OS and safety. DESTINY-Lung01 enrolled 181 patients at multiple sites, including Asia, Europe and North America. For more information about the trial, visit [ClinicalTrials.gov](https://ClinicalTrials.gov).

Full results from the *HER2* mutant cohort were [presented](#) at the 2021 ESMO Congress and simultaneously published in *The New England Journal of Medicine*. Updated results from both cohorts of DESTINY-Lung01 were [presented](#) at the 2022 ESMO Congress.

### **About DESTINY-CRC02**

DESTINY-CRC02 is a global, randomized, two arm, parallel, multicenter phase 2 trial evaluating the efficacy and safety of two doses (5.4 mg/kg or 6.4 mg/kg) of ENHERTU in patients with locally advanced, unresectable or metastatic *HER2* positive colorectal cancer of BRAF wild-type, or RAS wild-type and RAS mutant tumor types previously treated with standard therapy. The trial was conducted in two stages. In the first stage, patients (n=80) were randomized 1:1 to receive either 5.4 mg/kg or 6.4 mg/kg of ENHERTU. In the second stage, additional patients (n=42) were enrolled in the 5.4 mg/kg arm. The primary endpoint is confirmed ORR as assessed by blinded independent central review. Secondary endpoints include DOR, DCR, investigator-assessed confirmed ORR, clinical benefit ratio, PFS, OS and safety. DESTINY-CRC02 enrolled 122 patients at multiple sites in Asia, Europe and North America. For more information about the trial, visit [ClinicalTrials.gov](https://ClinicalTrials.gov).

Primary results from the DESTINY-CRC02 phase 2 trial were [presented](#) at the 2023 American Society of Clinical Oncology (ASCO) Annual Meeting.

### **About HER2 Expression in Solid Tumors**

*HER2* is a tyrosine kinase receptor growth-promoting protein expressed on the surface of various tissue cells throughout the body and is involved in normal cell growth.<sup>1,2</sup> In some cancers, *HER2* expression is amplified or the cells have activating mutations.<sup>1,3</sup> *HER2* protein overexpression may occur as a result of *HER2* gene amplification and is often associated with aggressive disease and poor prognosis.<sup>4</sup> While *HER2* directed therapies have been used to treat breast, gastric, lung and colorectal cancers, more research is needed evaluating their potential role in treating other *HER2* expressing solid tumor types.<sup>2,5,6</sup>

*HER2* is an emerging biomarker in solid tumor types including biliary tract, bladder, cervical, endometrial, ovarian and pancreatic cancers.<sup>3</sup> Testing is not routinely performed in these additional tumor types and as a result, available literature is limited. In these solid tumors, *HER2* positive expression, classified as IHC 3+, has been observed at rates from 1% to 28%.<sup>7,8</sup>

There is an unmet need for effective therapies for certain HER2 expressing solid tumors, particularly for those who have progressed on or are refractory to standard of care therapies as there are currently no approved HER2 directed therapies for these cancers.<sup>2,9</sup>

### **About ENHERTU**

ENHERTU (trastuzumab deruxtecan; fam-trastuzumab deruxtecan-nxki in the U.S. only) is a HER2 directed ADC. Designed using Daiichi Sankyo's proprietary DXd ADC technology, ENHERTU is the lead ADC in the oncology portfolio of Daiichi Sankyo and the most advanced program in AstraZeneca's ADC scientific platform. ENHERTU consists of a HER2 monoclonal antibody attached to a number of topoisomerase I inhibitor payloads (an exatecan derivative, DXd) via tetrapeptide-based cleavable linkers.

ENHERTU (5.4 mg/kg) is approved in more than 55 countries worldwide for the treatment of adult patients with unresectable or metastatic HER2 positive breast cancer who have received a (or one or more) prior anti-HER2-based regimen, either in the metastatic setting or in the neoadjuvant or adjuvant setting, and have developed disease recurrence during or within six months of completing therapy based on the results from the [DESTINY-Breast03](#) trial.

ENHERTU (5.4 mg/kg) is approved in more than 45 countries worldwide for the treatment of adult patients with unresectable or metastatic HER2 low (IHC 1+ or IHC 2+/*in-situ* hybridization (ISH)-) breast cancer who have received a prior systemic therapy in the metastatic setting or developed disease recurrence during or within six months of completing adjuvant chemotherapy based on the results from the [DESTINY-Breast04](#) trial.

ENHERTU (5.4 mg/kg) is approved in more than 30 countries worldwide for the treatment of adult patients with unresectable or metastatic NSCLC whose tumors have activating *HER2 (ERBB2)* mutations, as detected by a locally or regionally approved test, and who have received a prior systemic therapy based on the results from the [DESTINY-Lung02](#) trial. Continued approval in the U.S. for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

ENHERTU (6.4 mg/kg) is approved in more than 30 countries worldwide for the treatment of adult patients with locally advanced or metastatic HER2 positive gastric or gastroesophageal junction (GEJ) adenocarcinoma who have received a prior trastuzumab-based regimen based on the results from the [DESTINY-Gastric01](#) and/or [DESTINY-Gastric02](#) trials.

### **About the ENHERTU Clinical Development Program**

A comprehensive global clinical development program is underway evaluating the efficacy and safety of ENHERTU monotherapy across multiple HER2 targetable cancers. Trials in combination with other anticancer treatments, such as immunotherapy, also are underway.

### **About the Daiichi Sankyo and AstraZeneca Collaboration**

Daiichi Sankyo and AstraZeneca entered into a global collaboration to jointly develop and commercialize ENHERTU in [March 2019](#) and datopotamab deruxtecan (Dato-DXd) in [July 2020](#), except in Japan where Daiichi Sankyo maintains exclusive rights for each ADC. Daiichi Sankyo is responsible for the manufacturing and supply of ENHERTU and datopotamab deruxtecan.

### **About the DXd ADC Portfolio of Daiichi Sankyo**

The DXd ADC portfolio of Daiichi Sankyo currently consists of six ADCs in clinical development across multiple types of cancer. ENHERTU, a HER2 directed ADC, and datopotamab deruxtecan (Dato-DXd), a TROP2 directed ADC, are being jointly developed and commercialized globally with AstraZeneca.

Patritumab deruxtecan (HER3-DXd), a HER3 directed ADC, ifinatamab deruxtecan (I-DXd), a B7-H3 directed ADC, and raludotatug deruxtecan (R-DXd), a CDH6 directed ADC, are being jointly developed and commercialized globally with Merck & Co., Inc., Rahway, N.J. USA. DS-3939, a TA-MUC1 directed ADC, is being developed by Daiichi Sankyo.

Designed using Daiichi Sankyo's proprietary DXd ADC technology to target and deliver a cytotoxic payload inside cancer cells that express a specific cell surface antigen, each ADC consists of a monoclonal antibody attached to a number of topoisomerase I inhibitor payloads (an exatecan derivative, DXd) via tetrapeptide-based cleavable linkers.

Datopotamab deruxtecan, ifinatamab deruxtecan, patritumab deruxtecan, raludotatug deruxtecan and DS-3939 are investigational medicines that have not been approved for any indication in any country. Safety and efficacy have not been established.

## **ENHERTU U.S. Important Safety Information**

### **Indications**

ENHERTU is a HER2-directed antibody and topoisomerase inhibitor conjugate indicated for the treatment of adult patients with:

- Unresectable or metastatic HER2-positive breast cancer who have received a prior anti-HER2-based regimen either:
  - In the metastatic setting, or

- In the neoadjuvant or adjuvant setting and have developed disease recurrence during or within six months of completing therapy
- Unresectable or metastatic HER2-low (IHC 1+ or IHC 2+/ISH-) breast cancer, as determined by an FDA-approved test, who have received a prior chemotherapy in the metastatic setting or developed disease recurrence during or within 6 months of completing adjuvant chemotherapy
- Unresectable or metastatic non-small cell lung cancer (NSCLC) whose tumors have activating HER2 (ERBB2) mutations, as detected by an FDA-approved test, and who have received a prior systemic therapy

This indication is approved under accelerated approval based on objective response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

- Locally advanced or metastatic HER2-positive gastric or gastroesophageal junction adenocarcinoma who have received a prior trastuzumab-based regimen

**WARNING: INTERSTITIAL LUNG DISEASE and EMBRYO-FETAL TOXICITY**

- **Interstitial lung disease (ILD) and pneumonitis, including fatal cases, have been reported with ENHERTU. Monitor for and promptly investigate signs and symptoms including cough, dyspnea, fever, and other new or worsening respiratory symptoms. Permanently discontinue ENHERTU in all patients with Grade 2 or higher ILD/pneumonitis. Advise patients of the risk and to immediately report symptoms.**
- **Exposure to ENHERTU during pregnancy can cause embryo-fetal harm. Advise patients of these risks and the need for effective contraception.**

**Contraindications**

None.

**Warnings and Precautions**

**Interstitial Lung Disease / Pneumonitis**

Severe, life-threatening, or fatal interstitial lung disease (ILD), including pneumonitis, can occur in patients treated with ENHERTU. A higher incidence of Grade 1 and 2 ILD/pneumonitis has been observed in patients with moderate renal impairment. Advise patients to immediately report cough, dyspnea, fever, and/or any new or worsening respiratory symptoms. Monitor patients for signs and symptoms of ILD. Promptly investigate evidence of ILD. Evaluate patients with suspected ILD by radiographic imaging. Consider consultation with a pulmonologist. For asymptomatic ILD/pneumonitis (Grade 1), interrupt ENHERTU until resolved to Grade 0, then if resolved in  $\leq 28$  days from date of onset, maintain dose. If resolved in  $> 28$  days from date of onset, reduce dose one level. Consider corticosteroid treatment as soon as ILD/pneumonitis is suspected (e.g.,  $\geq 0.5$  mg/kg/day prednisolone or equivalent). For symptomatic ILD/pneumonitis (Grade 2 or greater), permanently discontinue ENHERTU. Promptly initiate systemic corticosteroid treatment as soon as ILD/pneumonitis is suspected (e.g.,  $\geq 1$  mg/kg/day prednisolone or equivalent) and continue for at least 14 days followed by gradual taper for at least 4 weeks.

Metastatic Breast Cancer and HER2-Mutant NSCLC (5.4 mg/kg)

In patients with metastatic breast cancer and HER2-mutant NSCLC treated with ENHERTU 5.4 mg/kg, ILD occurred in 12% of patients. Fatal outcomes due to ILD and/or pneumonitis occurred in 1.0% of patients treated with ENHERTU. Median time to first onset was 5 months (range: 0.9 to 23).

### Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

In patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg, ILD occurred in 10% of patients. Median time to first onset was 2.8 months (range: 1.2 to 21).

### **Neutropenia**

Severe neutropenia, including febrile neutropenia, can occur in patients treated with ENHERTU. Monitor complete blood counts prior to initiation of ENHERTU and prior to each dose, and as clinically indicated. For Grade 3 neutropenia (Absolute Neutrophil Count [ANC]  $<1.0$  to  $0.5 \times 10^9/L$ ), interrupt ENHERTU until resolved to Grade 2 or less, then maintain dose. For Grade 4 neutropenia (ANC  $<0.5 \times 10^9/L$ ), interrupt ENHERTU until resolved to Grade 2 or less, then reduce dose by one level. For febrile neutropenia (ANC  $<1.0 \times 10^9/L$  and temperature  $>38.3^\circ C$  or a sustained temperature of  $\geq 38^\circ C$  for more than 1 hour), interrupt ENHERTU until resolved, then reduce dose by one level.

### Metastatic Breast Cancer and HER2-Mutant NSCLC (5.4 mg/kg)

In patients with metastatic breast cancer and HER2-mutant NSCLC treated with ENHERTU 5.4 mg/kg, a decrease in neutrophil count was reported in 65% of patients. Sixteen percent had Grade 3 or 4 decreased neutrophil count. Median time to first onset of decreased neutrophil count was 22 days (range: 2 to 664). Febrile neutropenia was reported in 1.1% of patients.

### Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

In patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg, a decrease in neutrophil count was reported in 72% of patients. Fifty-one percent had Grade 3 or 4 decreased neutrophil count. Median time to first onset of decreased neutrophil count was 16 days (range: 4 to 187). Febrile neutropenia was reported in 4.8% of patients.

### **Left Ventricular Dysfunction**

Patients treated with ENHERTU may be at increased risk of developing left ventricular dysfunction. Left ventricular ejection fraction (LVEF) decrease has been observed with anti-HER2 therapies, including ENHERTU. Assess LVEF prior to initiation of ENHERTU and at regular intervals during treatment as clinically indicated. Manage LVEF decrease through treatment interruption. When LVEF is  $>45\%$  and absolute decrease from baseline is 10-20%, continue treatment with ENHERTU. When LVEF is 40-45% and absolute decrease from baseline is  $<10\%$ , continue treatment with ENHERTU and repeat LVEF assessment within 3 weeks. When LVEF is 40-45% and absolute decrease from baseline is 10-20%, interrupt ENHERTU and repeat LVEF assessment within 3 weeks. If LVEF has not recovered to within 10% from baseline, permanently discontinue ENHERTU. If LVEF recovers to within 10% from baseline, resume treatment with ENHERTU at the same dose. When LVEF is  $<40\%$  or absolute decrease from baseline is  $>20\%$ , interrupt ENHERTU and repeat LVEF assessment within 3 weeks. If LVEF of  $<40\%$  or absolute decrease from baseline of  $>20\%$  is confirmed, permanently discontinue ENHERTU. Permanently discontinue ENHERTU in patients with symptomatic congestive heart failure. Treatment with ENHERTU has not been studied in patients with a history of clinically significant cardiac disease or LVEF  $<50\%$  prior to initiation of treatment.

### Metastatic Breast Cancer and HER2-Mutant NSCLC (5.4 mg/kg)

In patients with metastatic breast cancer and HER2-mutant NSCLC treated with ENHERTU 5.4 mg/kg, LVEF decrease was reported in 3.6% of patients, of which 0.4% were Grade 3.

### Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

In patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg, no clinical adverse events of heart failure were reported; however, on echocardiography, 8% were found to have asymptomatic Grade 2 decrease in LVEF.

### **Embryo-Fetal Toxicity**

ENHERTU can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risks to a fetus. Verify the pregnancy status of females of reproductive potential prior to the initiation of

ENHERTU. Advise females of reproductive potential to use effective contraception during treatment and for 7 months after the last dose of ENHERTU. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ENHERTU and for 4 months after the last dose of ENHERTU.

### **Additional Dose Modifications**

#### **Thrombocytopenia**

For Grade 3 thrombocytopenia (platelets  $<50$  to  $25 \times 10^9/L$ ) interrupt ENHERTU until resolved to Grade 1 or less, then maintain dose. For Grade 4 thrombocytopenia (platelets  $<25 \times 10^9/L$ ) interrupt ENHERTU until resolved to Grade 1 or less, then reduce dose by one level.

### **Adverse Reactions**

#### Metastatic Breast Cancer and HER2-Mutant NSCLC (5.4 mg/kg)

The pooled safety population reflects exposure to ENHERTU 5.4 mg/kg intravenously every 3 weeks in 984 patients in Study DS8201-A-J101 (NCT02564900), DESTINY-Breast01, DESTINY-Breast03, DESTINY-Breast04, and DESTINY-Lung02. Among these patients 65% were exposed for  $>6$  months and 39% were exposed for  $>1$  year. In this pooled safety population, the most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were nausea (76%), decreased white blood cell count (71%), decreased hemoglobin (66%), decreased neutrophil count (65%), decreased lymphocyte count (55%), fatigue (54%), decreased platelet count (47%), increased aspartate aminotransferase (48%), vomiting (44%), increased alanine aminotransferase (42%), alopecia (39%), increased blood alkaline phosphatase (39%), constipation (34%), musculoskeletal pain (32%), decreased appetite (32%), hypokalemia (28%), diarrhea (28%), and respiratory infection (24%).

#### HER2-Positive Metastatic Breast Cancer

##### *DESTINY-Breast03*

The safety of ENHERTU was evaluated in 257 patients with unresectable or metastatic HER2-positive breast cancer who received at least one dose of ENHERTU 5.4 mg/kg intravenously every three weeks in DESTINY-Breast03. The median duration of treatment was 14 months (range: 0.7 to 30).

Serious adverse reactions occurred in 19% of patients receiving ENHERTU. Serious adverse reactions in  $>1\%$  of patients who received ENHERTU were vomiting, interstitial lung disease, pneumonia, pyrexia, and urinary tract infection. Fatalities due to adverse reactions occurred in 0.8% of patients including COVID-19 and sudden death (one patient each).

ENHERTU was permanently discontinued in 14% of patients, of which ILD/pneumonitis accounted for 8%. Dose interruptions due to adverse reactions occurred in 44% of patients treated with ENHERTU. The most frequent adverse reactions ( $>2\%$ ) associated with dose interruption were neutropenia, leukopenia, anemia, thrombocytopenia, pneumonia, nausea, fatigue, and ILD/pneumonitis. Dose reductions occurred in 21% of patients treated with ENHERTU. The most frequent adverse reactions ( $>2\%$ ) associated with dose reduction were nausea, neutropenia, and fatigue.

The most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were nausea (76%), decreased white blood cell count (74%), decreased neutrophil count (70%), increased aspartate aminotransferase (67%), decreased hemoglobin (64%), decreased lymphocyte count (55%), increased alanine aminotransferase (53%), decreased platelet count (52%), fatigue (49%), vomiting (49%), increased blood alkaline phosphatase (49%), alopecia (37%), hypokalemia (35%), constipation (34%), musculoskeletal pain (31%), diarrhea (29%), decreased appetite (29%), respiratory infection (22%), headache (22%), abdominal pain (21%), increased blood bilirubin (20%), and stomatitis (20%).

#### HER2-Low Metastatic Breast Cancer

##### *DESTINY-Breast04*

The safety of ENHERTU was evaluated in 371 patients with unresectable or metastatic HER2-low (IHC 1+ or IHC 2+/ISH-) breast cancer who received ENHERTU 5.4 mg/kg intravenously every 3 weeks in



DESTINY-Breast04. The median duration of treatment was 8 months (range: 0.2 to 33) for patients who received ENHERTU.

Serious adverse reactions occurred in 28% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were ILD/pneumonitis, pneumonia, dyspnea, musculoskeletal pain, sepsis, anemia, febrile neutropenia, hypercalcemia, nausea, pyrexia, and vomiting. Fatalities due to adverse reactions occurred in 4% of patients including ILD/pneumonitis (3 patients); sepsis (2 patients); and ischemic colitis, disseminated intravascular coagulation, dyspnea, febrile neutropenia, general physical health deterioration, pleural effusion, and respiratory failure (1 patient each).

ENHERTU was permanently discontinued in 16% of patients, of which ILD/pneumonitis accounted for 8%. Dose interruptions due to adverse reactions occurred in 39% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose interruption were neutropenia, fatigue, anemia, leukopenia, COVID-19, ILD/pneumonitis, increased transaminases, and hyperbilirubinemia. Dose reductions occurred in 23% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were fatigue, nausea, thrombocytopenia, and neutropenia.

The most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were nausea (76%), decreased white blood cell count (70%), decreased hemoglobin (64%), decreased neutrophil count (64%), decreased lymphocyte count (55%), fatigue (54%), decreased platelet count (44%), alopecia (40%), vomiting (40%), increased aspartate aminotransferase (38%), increased alanine aminotransferase (36%), constipation (34%), increased blood alkaline phosphatase (34%), decreased appetite (32%), musculoskeletal pain (32%), diarrhea (27%), and hypokalemia (25%).

#### Unresectable or Metastatic HER2-Mutant NSCLC (5.4 mg/kg)

DESTINY-Lung02 evaluated two dose levels (5.4 mg/kg [n=101] and 6.4 mg/kg [n=50]); however, only the results for the recommended dose of 5.4 mg/kg intravenously every 3 weeks are described below due to increased toxicity observed with the higher dose in patients with NSCLC, including ILD/pneumonitis.

The safety of ENHERTU was evaluated in 101 patients with unresectable or metastatic HER2-mutant NSCLC who received ENHERTU 5.4 mg/kg intravenously every three weeks in DESTINY-Lung02. Nineteen percent of patients were exposed for >6 months.

Serious adverse reactions occurred in 30% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were ILD/pneumonitis, thrombocytopenia, dyspnea, nausea, pleural effusion, and increased troponin I. Fatality occurred in 1 patient with suspected ILD/pneumonitis (1%).

ENHERTU was permanently discontinued in 8% of patients. Adverse reactions which resulted in permanent discontinuation of ENHERTU were ILD/pneumonitis, diarrhea, hypokalemia, hypomagnesemia, myocarditis, and vomiting. Dose interruptions of ENHERTU due to adverse reactions occurred in 23% of patients. Adverse reactions which required dose interruption (>2%) included neutropenia and ILD/pneumonitis. Dose reductions due to an adverse reaction occurred in 11% of patients.

The most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were nausea (61%), decreased white blood cell count (60%), decreased hemoglobin (58%), decreased neutrophil count (52%), decreased lymphocyte count (43%), decreased platelet count (40%), decreased albumin (39%), increased aspartate aminotransferase (35%), increased alanine aminotransferase (34%), fatigue (32%), constipation (31%), decreased appetite (30%), vomiting (26%), increased alkaline phosphatase (22%), and alopecia (21%).

### Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

The safety of ENHERTU was evaluated in 187 patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma in DESTINY-Gastric01. Patients intravenously received at least one dose of either ENHERTU (N=125) 6.4 mg/kg every 3 weeks or either irinotecan (N=55) 150 mg/m<sup>2</sup> biweekly or paclitaxel (N=7) 80 mg/m<sup>2</sup> weekly for 3 weeks. The median duration of treatment was 4.6 months (range: 0.7 to 22.3) for patients who received ENHERTU.

Serious adverse reactions occurred in 44% of patients receiving ENHERTU 6.4 mg/kg. Serious adverse reactions in >2% of patients who received ENHERTU were decreased appetite, ILD, anemia, dehydration, pneumonia, cholestatic jaundice, pyrexia, and tumor hemorrhage. Fatalities due to adverse reactions occurred in 2.4% of patients: disseminated intravascular coagulation, large intestine perforation, and pneumonia occurred in one patient each (0.8%).

ENHERTU was permanently discontinued in 15% of patients, of which ILD accounted for 6%. Dose interruptions due to adverse reactions occurred in 62% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose interruption were neutropenia, anemia, decreased appetite, leukopenia, fatigue, thrombocytopenia, ILD, pneumonia, lymphopenia, upper respiratory tract infection, diarrhea, and hypokalemia. Dose reductions occurred in 32% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were neutropenia, decreased appetite, fatigue, nausea, and febrile neutropenia.

The most common ( $\geq 20\%$ ) adverse reactions, including laboratory abnormalities, were decreased hemoglobin (75%), decreased white blood cell count (74%), decreased neutrophil count (72%), decreased lymphocyte count (70%), decreased platelet count (68%), nausea (63%), decreased appetite (60%), increased aspartate aminotransferase (58%), fatigue (55%), increased blood alkaline phosphatase (54%), increased alanine aminotransferase (47%), diarrhea (32%), hypokalemia (30%), vomiting (26%), constipation (24%), increased blood bilirubin (24%), pyrexia (24%), and alopecia (22%).

### **Use in Specific Populations**

- **Pregnancy:** ENHERTU can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risks to a fetus. There are clinical considerations if ENHERTU is used in pregnant women, or if a patient becomes pregnant within 7 months after the last dose of ENHERTU.
- **Lactation:** There are no data regarding the presence of ENHERTU in human milk, the effects on the breastfed child, or the effects on milk production. Because of the potential for serious adverse reactions in a breastfed child, advise women not to breastfeed during treatment with ENHERTU and for 7 months after the last dose.
- **Females and Males of Reproductive Potential:** Pregnancy testing: Verify pregnancy status of females of reproductive potential prior to initiation of ENHERTU. Contraception: *Females:* ENHERTU can cause fetal harm when administered to a pregnant woman. Advise females of reproductive potential to use effective contraception during treatment with ENHERTU and for 7 months after the last dose. *Males:* Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ENHERTU and for 4 months after the last dose. Infertility: ENHERTU may impair male reproductive function and fertility.
- **Pediatric Use:** Safety and effectiveness of ENHERTU have not been established in pediatric patients.
- **Geriatric Use:** Of the 883 patients with breast cancer treated with ENHERTU 5.4 mg/kg, 22% were  $\geq 65$  years and 3.6% were  $\geq 75$  years. No overall differences in efficacy within clinical studies were observed between patients  $\geq 65$  years of age compared to younger patients. There was a higher incidence of Grade 3-4 adverse reactions observed in patients aged  $\geq 65$  years (60%) as compared to younger patients (48%). Of the 101 patients with unresectable or metastatic HER2-mutant NSCLC treated with ENHERTU 5.4 mg/kg, 40% were  $\geq 65$  years and 8% were  $\geq 75$  years. No overall differences in efficacy or safety were observed between patients  $\geq 65$  years of age compared to younger patients. Of the 125 patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg in DESTINY-Gastric01, 56% were  $\geq 65$  years and 14% were  $\geq 75$  years. No overall differences in efficacy or safety were observed between patients  $\geq 65$  years of age compared to younger patients.

- **Renal Impairment:** A higher incidence of Grade 1 and 2 ILD/pneumonitis has been observed in patients with moderate renal impairment. Monitor patients with moderate renal impairment more frequently. The recommended dosage of ENHERTU has not been established for patients with severe renal impairment (CLCr <30 mL/min).
- **Hepatic Impairment:** In patients with moderate hepatic impairment, due to potentially increased exposure, closely monitor for increased toxicities related to the topoisomerase inhibitor. The recommended dosage of ENHERTU has not been established for patients with severe hepatic impairment (total bilirubin >3 times ULN and any AST).

**To report SUSPECTED ADVERSE REACTIONS, contact Daiichi Sankyo, Inc. at 1-877-437-7763 or FDA at 1-800-FDA-1088 or [fda.gov/medwatch](http://fda.gov/medwatch).**

**Please see accompanying full [Prescribing Information](#), including [Boxed WARNINGS](#), and [Medication Guide](#).**

### About Daiichi Sankyo

Daiichi Sankyo is an innovative global healthcare company contributing to the sustainable development of society that discovers, develops and delivers new standards of care to enrich the quality of life around the world. With more than 120 years of experience, Daiichi Sankyo leverages its world-class science and technology to create new modalities and innovative medicines for people with cancer, cardiovascular and other diseases with high unmet medical need. For more information, please visit [www.daiichisankyo.com](http://www.daiichisankyo.com).

### Media Contacts:

#### Global/US:

Jennifer Brennan  
Daiichi Sankyo, Inc.  
[jbrennan2@dsi.com](mailto:jbrennan2@dsi.com)  
+1 908 900 3183 (mobile)

#### Japan:

Koji Ogiwara  
Daiichi Sankyo Co., Ltd.  
[ogiwara.koji.ay@daiichisankyo.co.jp](mailto:ogiwara.koji.ay@daiichisankyo.co.jp)  
+81 3 6225 1126 (office)

#### Investor Relations Contact:

[DaiichiSankyoIR@daiichisankyo.co.jp](mailto:DaiichiSankyoIR@daiichisankyo.co.jp)

---

#### References:

- <sup>1</sup> ASCO. [Breast Cancer](#). Accessed January 2024.
- <sup>2</sup> Iqbal N, et al. *Mol Biol Int*. 2014;852748.
- <sup>3</sup> Omar N, et al. *Pathogenesis*. 2015;2(3):1-9.
- <sup>4</sup> Pillai R, et al. *Cancer*. 2017;1;123(21): 4099-4105.
- <sup>5</sup> National Cancer Institute. [Enhertu Marks First Targeted Therapy for HER2-Mutant Lung Cancer](#). Accessed January 2024.
- <sup>6</sup> Siena S, et al. *Ann Oncol*. 2018 May; 29(5):1108–1119.
- <sup>7</sup> Yan M, et al. *Cancer Metastasis Rev*. 2015;34(1):157–164.
- <sup>8</sup> Buza N, et al. *Modern Pathology*. 2013 Dec;26(12):1605-12.
- <sup>9</sup> Meric-Bernstam F, et al. *J Clin Oncol*. 2021; 39:15: 3004-3004.