

Enhertu® Wins 2026 International Prix Galien Award for Best Biotechnology Product

- Second Prix Galien Award for Daiichi Sankyo and AstraZeneca's Enhertu following receipt of the Prix Galien USA Award in 2024

Tokyo – (October 30, 2026) – Enhertu® (trastuzumab deruxtecan) has been awarded the 2026 International Prix Galien Award for Best Biotechnology Product, a recognition awarded every two years by The Galien Foundation. This is the second award granted to Enhertu, following receipt of the Prix Galien USA Award for Best Biotechnology Product in 2024.

The Prix Galien is among the global health innovation industry's most renowned honors, recognizing outstanding biomedical and medical technology product achievements that improve the human condition, and is considered the industry's equivalent of the Nobel Prize. Medicines that have been recognized with a National Prix Galien Award within the prior two years are eligible for the International Prix Galien Award. Enhertu was one of six nominees for the 2026 International Prix Galien Award for Best Biotechnology Product. Winners are selected by an international committee comprised of chairs from each of the member nations, including Africa, Belgium/Luxembourg, Bridges, Canada, China, France, Germany, Greece, India, Italy, Japan, Korea, Poland, Switzerland, the Netherlands, the UK and the USA. The award was presented during a ceremony at the American Museum of Natural History in New York on Thursday, October 29, 2026.

Enhertu is a specifically engineered HER2 directed DXd antibody drug conjugate (ADC) discovered by Daiichi Sankyo (TSE: 4568) and being jointly developed and commercialized by Daiichi Sankyo and AstraZeneca (LSE/STO/NYSE: AZN).

"We are honored to receive this second Prix Galien award, which recognizes the impressive advances Enhertu has made possible for certain patients with cancer around the world. Since its first U.S. approval in 2019, Enhertu has transformed the treatment and classification of metastatic breast cancer by pioneering the establishment of HER2 low and HER2 ultralow breast cancer as clinically distinct patient segments," said Hiroyuki Okuzawa, President and CEO of Daiichi Sankyo. "Enhertu also has delivered a series of additional practice-changing results that have reshaped standards of care across early stage and metastatic settings of HER2 positive breast cancer and HER2 positive metastatic gastric cancer, while also becoming the first HER2 directed tumor agnostic medicine for HER2 positive solid tumors. We are extremely proud that scientists at Daiichi Sankyo invented the DXd antibody drug conjugate technology behind Enhertu, and we look to continue this important work through the development of additional breakthrough medicines for cancer."

The Galien Foundation fosters, recognizes and rewards excellence in scientific innovation to improve human health. Its vision is to serve as a catalyst for the development of the next generation of treatments and technologies that will transform medical practice and save lives. Founded in France in 1970 by pharmacist Roland Mehl in honor of Galen, the father of medical science and modern pharmacology, the Prix Galien is widely regarded as the equivalent of the Nobel Prize in biopharmaceutical research.

About Enhertu

Enhertu (trastuzumab deruxtecan; fam-trastuzumab deruxtecan-nxki in the U.S. only) is a HER2 directed ADC. Designed using the proprietary DXd ADC Technology of Daiichi Sankyo, Enhertu is the lead ADC in the oncology portfolio of Daiichi Sankyo and the most advanced program in AstraZeneca's ADC scientific platform. Enhertu consists of a HER2 monoclonal antibody attached to a number of topoisomerase I inhibitor payloads (an exatecan derivative, DXd) via tetrapeptide-based cleavable linkers.

Enhertu (5.4 mg/kg) followed by THP is approved in more than five countries/regions worldwide as a neoadjuvant treatment for adult patients with HER2 positive (IHC 3+ or ISH+) stage 2 or stage 3 breast cancer based on the results from the [DESTINY-Breast11](#) trial.

Enhertu (5.4 mg/kg) is approved in more than 35 countries/regions worldwide for the adjuvant treatment of adult patients with HER2 positive breast cancer who have residual invasive disease following neoadjuvant trastuzumab (with or without pertuzumab) and taxane-based treatment based on the [DESTINY-Breast05](#) trial.

Enhertu (5.4 mg/kg) in combination with pertuzumab is approved in more than 40 countries/regions worldwide as a first-line treatment for adult patients with unresectable or metastatic HER2 positive (IHC 3+ or ISH+) breast cancer, as determined by a locally or regionally approved test, based on the results from the [DESTINY-Breast09](#) trial.

Enhertu (5.4 mg/kg) is approved in more than 100 countries/regions worldwide for the treatment of adult patients with unresectable or metastatic HER2 positive (IHC 3+ or ISH+) breast cancer who have received a prior anti-HER2-based regimen, either in the metastatic setting or in the neoadjuvant or adjuvant setting, and have developed disease recurrence during or within six months of completing therapy based on the results from the [DESTINY-Breast03](#) trial.

Enhertu (5.4 mg/kg) is approved in more than 75 countries/regions worldwide for the treatment of adult patients with unresectable or metastatic hormone receptor (HR) positive, HER2 low (IHC 1+ or IHC 2+/ISH-) or HER2 ultralow (IHC 0 with membrane staining) breast cancer, as determined by a locally or regionally approved test, that have progressed on one or more endocrine therapies in the metastatic setting based on the results from the [DESTINY-Breast06](#) trial.

Enhertu (5.4 mg/kg) is approved in more than 100 countries/regions worldwide for the treatment of adult patients with unresectable or metastatic HER2 low (IHC 1+ or IHC 2+/ISH-) breast cancer who have received a prior systemic therapy in the metastatic setting or developed disease recurrence during or within six months of completing adjuvant chemotherapy based on the results from the [DESTINY-Breast04](#) trial.

Enhertu (5.4 mg/kg) is approved in more than 80 countries/regions worldwide for the treatment of adult patients with unresectable or metastatic NSCLC whose tumors have activating *HER2* (*ERBB2*) mutations, as detected by a locally or regionally approved test, and who have received a prior systemic therapy based on the results from the [DESTINY-Lung02](#) and/or [DESTINY-Lung05](#) trials. Continued approval in China and the U.S. for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

Enhertu (6.4 mg/kg) is approved in more than 90 countries/regions worldwide for the treatment of adult patients with locally advanced or metastatic HER2 positive (IHC 3+ or IHC 2+/ISH+) gastric or gastroesophageal junction (GEJ) adenocarcinoma who have received a prior trastuzumab-based regimen based on the results from the [DESTINY-Gastric01](#), [DESTINY-Gastric02](#) and/or [DESTINY-Gastric04](#) trials.

Enhertu (5.4 mg/kg) is approved in more than 50 countries/regions worldwide for the treatment of adult patients with unresectable or metastatic HER2 positive (IHC 3+) solid tumors who have received prior treatment and have no satisfactory alternative treatment options based on efficacy results from the [DESTINY-PanTumor02](#), [DESTINY-Lung01](#), [DESTINY-CRC02](#), [DESTINY-PanTumor03](#) and/or [HERALD](#) trials. Continued approval in China and the U.S. for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

About the Enhertu Clinical Development Program

A comprehensive global clinical development program is underway evaluating the efficacy and safety of Enhertu as a monotherapy or in combination or sequentially with other cancer medicines across multiple HER2 targetable cancers.

About the Daiichi Sankyo and AstraZeneca Collaboration

Daiichi Sankyo and AstraZeneca entered into a global collaboration to jointly develop and commercialize Enhertu in [March 2019](#) and Datroway® in [July 2020](#), except in Japan where Daiichi Sankyo maintains exclusive rights for each ADC. Daiichi Sankyo is responsible for the manufacturing and supply of Enhertu and Datroway.

Enhertu U.S. Indications and Important Safety Information

Indications

ENHERTU is a HER2-directed antibody and topoisomerase inhibitor conjugate indicated for:

- HER2-Positive Early Breast Cancer
 - As neoadjuvant treatment of adult patients with HER2-positive (IHC 3+ or ISH+) Stage II or III breast cancer, as determined by an FDA-authorized test followed by a taxane, trastuzumab, and pertuzumab (THP)
 - As monotherapy in adult patients with HER2-positive (IHC 3+ or ISH+) breast cancer who have residual invasive disease after neoadjuvant HER2-targeted treatment
- HER2-Positive Metastatic Breast Cancer
 - In combination with pertuzumab as first-line treatment of adult patients with unresectable or metastatic HER2-positive (IHC 3+ or ISH+) breast cancer, as determined by an FDA-authorized test
 - As monotherapy for the treatment of adult patients with unresectable or metastatic HER2-positive (IHC 3+ or ISH+) breast cancer who have received a prior anti-HER2-based regimen either in the metastatic setting, or, in the neoadjuvant or adjuvant setting and have developed disease recurrence during or within six months of completing therapy
- HER2-Low and HER2-Ultralow Metastatic Breast Cancer
 - As monotherapy for the treatment of adult patients with unresectable or metastatic hormone receptor (HR)-positive, HER2-low (IHC 1+ or IHC 2+/ISH-) or HER2-ultralow (IHC 0 with membrane staining) breast cancer, as determined by an FDA-authorized test, that has progressed on one or more endocrine therapies in the metastatic setting
 - As monotherapy for the treatment of adult patients with unresectable or metastatic HER2-low (IHC 1+ or IHC 2+/ISH-) breast cancer, as determined by an FDA-authorized test, who have received a prior chemotherapy in the metastatic setting or developed disease recurrence during or within 6 months of completing adjuvant chemotherapy
- HER2-Mutant Unresectable or Metastatic Non-Small Cell Lung Cancer (NSCLC)
 - As monotherapy for the treatment of adult patients with unresectable or metastatic NSCLC whose tumors have activating HER2 (ERBB2) mutations, as detected by an FDA-authorized test, and who have received a prior systemic therapy

This indication is approved under accelerated approval based on objective response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

- HER2-Positive Locally Advanced or Metastatic Gastric Cancer
 - As monotherapy for the treatment of adult patients with locally advanced or metastatic HER2-positive (IHC 3+ or IHC 2+/ISH positive) gastric or gastroesophageal junction (GEJ) adenocarcinoma who have received a prior trastuzumab-based regimen

- HER2-Positive (IHC 3+) Unresectable or Metastatic Solid Tumors

- As monotherapy for the treatment of adult patients with unresectable or metastatic HER2-positive (IHC 3+) solid tumors who have received prior systemic treatment and have no satisfactory alternative treatment options

This indication is approved under accelerated approval based on objective response rate and duration of response. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial.

Important Safety Information

WARNING: INTERSTITIAL LUNG DISEASE and EMBRYO-FETAL TOXICITY

- **Interstitial lung disease (ILD) and pneumonitis, including severe, life-threatening, and fatal cases, have been reported with ENHERTU. Monitor for and promptly investigate signs and symptoms including cough, dyspnea, fever, and other new or worsening respiratory symptoms. Permanently discontinue ENHERTU in all patients with Grade 2 or higher ILD/pneumonitis. Advise patients of the risk and to immediately report symptoms.**
- **Exposure to ENHERTU during pregnancy can cause embryo-fetal harm. Advise patients of these risks and the need for effective contraception.**

Contraindications

None.

Warnings and Precautions

Interstitial Lung Disease / Pneumonitis

Severe, life-threatening, or fatal interstitial lung disease (ILD), including pneumonitis, can occur in patients treated with ENHERTU. A higher incidence of Grade 1 and 2 ILD/pneumonitis has been observed in patients with moderate renal impairment. Advise patients to immediately report cough, dyspnea, fever, and/or any new or worsening respiratory symptoms. Monitor patients for signs and symptoms of ILD. Promptly investigate evidence of ILD. Evaluate patients with suspected ILD by radiographic imaging. Consider consultation with a pulmonologist. For asymptomatic ILD/pneumonitis (Grade 1), interrupt ENHERTU until resolved to Grade 0, then if resolved in ≤28 days from date of onset, maintain dose. If resolved in >28 days from date of onset, reduce dose 1 level. Consider corticosteroid treatment as soon as ILD/pneumonitis is suspected (e.g., ≥0.5 mg/kg/day prednisolone or equivalent). For symptomatic ILD/pneumonitis (Grade 2 or greater), permanently discontinue ENHERTU. Promptly initiate systemic corticosteroid treatment as soon as ILD/pneumonitis is suspected (e.g., ≥1 mg/kg/day prednisolone or equivalent) and continue for at least 14 days followed by gradual taper for at least 4 weeks.

HER2-Positive, HER2-Low, and HER2-Ultralow Breast Cancer, HER2-Mutant NSCLC, and Solid Tumors (Including IHC 3+) (5.4 mg/kg)

ENHERTU as Monotherapy

In patients treated with ENHERTU 5.4 mg/kg, ILD occurred in 12% of patients. Median time to first onset was 5.5 months (range: 0.9 to 31.5). Fatal outcomes due to ILD and/or pneumonitis occurred in 0.9% of patients treated with ENHERTU.

ENHERTU in Combination with Pertuzumab

In patients treated with ENHERTU 5.4 mg/kg in combination with pertuzumab (N=431), ILD occurred in 12% of patients. Median time to first onset was 8.0 months (range: 0.6 to 33.8). Fatal outcomes due to ILD and/or pneumonitis occurred in 0.5% of patients treated with ENHERTU in combination with pertuzumab.

ENHERTU followed by THP

In patients treated with ENHERTU 5.4 mg/kg followed by THP in DESTINY-Breast11, ILD occurred in 4.4% of patients. Median time to first onset was 2.7 months (range: 1.1 to 6.0). Fatal outcomes due to ILD and/or pneumonitis occurred in 1 patient (0.3%) treated with ENHERTU followed by THP.

HER2-Positive Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

In patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg, ILD occurred in 10% of patients. Median time to first onset was 2.8 months (range: 1.2 to 21).

Neutropenia

Severe neutropenia, including febrile neutropenia, can occur in patients treated with ENHERTU. Monitor complete blood counts prior to initiation of ENHERTU and prior to each dose, and as clinically indicated. For Grade 3 neutropenia (Absolute Neutrophil Count [ANC] <1.0 to $0.5 \times 10^9/L$), interrupt ENHERTU until resolved to Grade 2 or less, then maintain dose. For Grade 4 neutropenia (ANC $<0.5 \times 10^9/L$), interrupt ENHERTU until resolved to Grade 2 or less, then reduce dose by 1 level. For febrile neutropenia (ANC $<1.0 \times 10^9/L$ and temperature $>38.3^\circ C$ or a sustained temperature of $\geq 38^\circ C$ for more than 1 hour), interrupt ENHERTU until resolved, then reduce dose by 1 level.

HER2-Positive, HER2-Low, and HER2-Ultralow Breast Cancer, HER2-Mutant NSCLC, and Solid Tumors (Including IHC 3+) (5.4 mg/kg)

ENHERTU as Monotherapy

In patients treated with ENHERTU 5.4 mg/kg, a decrease in neutrophil count was reported in 65% of patients. Nineteen percent had Grade 3 or 4 decreased neutrophil count. Median time to first onset of decreased neutrophil count was 22 days (range: 2 to 939). Febrile neutropenia was reported in 1% of patients.

ENHERTU in Combination with Pertuzumab

In patients treated with ENHERTU 5.4 mg/kg in combination with pertuzumab (N=431), decreased neutrophil count occurred in 79% of patients. Median time to first onset was 22 days (range: 5 to 994). Twenty-nine percent had Grade 3 or 4 decreased neutrophil count. Febrile neutropenia was reported in 2.6% of patients.

ENHERTU followed by THP

In patients treated with ENHERTU 5.4 mg/kg followed by THP in DESTINY-Breast11, a decrease in neutrophil count was reported in 58% of patients. Seventeen percent had Grade 3 or 4 decreased neutrophil count. Median time to first onset of decreased neutrophil count was 42 days (range: 11 to 165). Febrile neutropenia was reported in 0.9% of patients.

HER2-Positive Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

In patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg, a decrease in neutrophil count was reported in 72% of patients. Fifty-one percent had Grade 3 or 4 decreased neutrophil count. Median time to first onset of decreased neutrophil count was 16 days (range: 4 to 187). Febrile neutropenia was reported in 4.8% of patients.

Left Ventricular Dysfunction

Patients treated with ENHERTU may be at increased risk of developing left ventricular dysfunction. Left ventricular dysfunction (LVD) has been observed with anti-HER2 therapies, including ENHERTU. Assess left ventricular ejection fraction (LVEF) prior to initiation of ENHERTU and at regular intervals during treatment as clinically indicated. Manage LVD through treatment interruption. When LVEF is $>45\%$ and absolute decrease from baseline is 10-20%, continue treatment with ENHERTU. When LVEF is 40-45% and absolute decrease from baseline is $<10\%$, continue treatment with ENHERTU and repeat LVEF assessment within 3 weeks. When LVEF is 40-45% and absolute decrease from baseline is 10-20%, interrupt ENHERTU and repeat LVEF assessment within 3 weeks. If LVEF has not recovered to within 10% from baseline, permanently discontinue ENHERTU. If LVEF recovers to within 10% from baseline, resume treatment with ENHERTU at the same dose. When LVEF is $<40\%$ or absolute decrease from baseline is $>20\%$, interrupt ENHERTU and repeat LVEF assessment within 3 weeks. If LVEF of $<40\%$ or absolute decrease from baseline of $>20\%$ is confirmed, permanently discontinue ENHERTU. Permanently discontinue ENHERTU in patients with symptomatic congestive heart failure. Treatment with ENHERTU has not been studied in patients with a history of clinically significant cardiac disease or LVEF $<50\%$ prior to initiation of treatment.

HER2-Positive, HER2-Low, and HER2-Ultralow Breast Cancer, HER2-Mutant NSCLC, and Solid Tumors (Including IHC 3+) (5.4 mg/kg)

ENHERTU as Monotherapy

In patients treated with ENHERTU 5.4 mg/kg, LVD was reported in 4.6% of patients, of which 0.6% were Grade 3 or 4.

ENHERTU in Combination with Pertuzumab

In patients treated with ENHERTU 5.4 mg/kg in combination with pertuzumab (N=431), LVEF decrease was reported in 11% of patients, of which 2.1% were Grade 3 or 4.

ENHERTU followed by THP

In patients treated with ENHERTU 5.4 mg/kg followed by THP in DESTINY-Breast11, LVD was reported in 1.3% of patients, of which 0.3% were Grade 3.

HER2-Positive Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

In patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg, no clinical adverse events of heart failure were reported; however, on echocardiography, 8% were found to have asymptomatic Grade 2 decrease in LVEF.

Embryo-Fetal Toxicity

ENHERTU can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risks to a fetus. Verify the pregnancy status of females of reproductive potential prior to the initiation of ENHERTU. Advise females of reproductive potential to use effective contraception during treatment and for 7 months after the last dose of ENHERTU. Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ENHERTU and for 4 months after the last dose of ENHERTU.

Additional Dose Modifications

Thrombocytopenia

For Grade 3 thrombocytopenia (platelets <50 to $25 \times 10^9/L$) interrupt ENHERTU until resolved to Grade 1 or less, then maintain dose. For Grade 4 thrombocytopenia (platelets $<25 \times 10^9/L$) interrupt ENHERTU until resolved to Grade 1 or less, then reduce dose by 1 level.

Radiotherapy-Related Pneumonitis and Skin Injury in the Post-Neoadjuvant Early Breast Cancer Setting

Among patients receiving sequential or concurrent adjuvant radiotherapy in DESTINY-Breast05, radiation pneumonitis occurred in 31% of patients receiving ENHERTU 5.4 mg/kg (N=757) and 31% of patients receiving ado-trastuzumab emtansine (T-DM1) 3.6 mg/kg (N=750), as reported by the investigator. All cases of patients receiving ENHERTU and adjuvant radiotherapy (ART) were either Grade 1 (27%) or Grade 2 (4.9%). A higher incidence of radiation pneumonitis was reported for patients who received sequential vs concurrent ART: 34% with sequential ART and 29% with concurrent ART. For patients treated with ENHERTU, median time to onset of first event of radiation pneumonitis was 4.1 months (range: 1.3 to 11.6). Among 39% of patients with recovery of first event of radiation pneumonitis, median duration was 3.5 months (range: <0.1 to 33.8). Among patients receiving ART and ENHERTU, radiation skin injury occurred in 20% of all patients, of which 1.6% were Grade 3 and none were Grade 4. For Grade 1 (asymptomatic) radiotherapy-related pneumonitis, continue treatment with ENHERTU. For Grade 2 (symptomatic), interrupt ENHERTU until recovered to baseline or Grade ≤ 1 . For Grade ≥ 3 (symptomatic, severe, or life-threatening), permanently discontinue ENHERTU. Manage Grade ≥ 2 per local institutional guidelines.

Adverse Reactions

HER2-Positive, HER2-Low, and HER2-Ultralow Breast Cancer, HER2-Mutant NSCLC, and Solid Tumors (Including IHC 3+) (5.4 mg/kg)

ENHERTU as Monotherapy

The pooled safety population reflects exposure to ENHERTU 5.4 mg/kg intravenously every 3 weeks in 2233 patients in Study DS8201-A-J101 (NCT02564900), DESTINY-Breast01, DESTINY-Breast02, DESTINY-Breast03, DESTINY-Breast04, DESTINY-Breast06, DESTINY-Lung01, DESTINY-Lung02, DESTINY-CRC02, and DESTINY-PanTumor02. Among these patients, 67% were exposed for >6 months and 39% were exposed for >1 year. In this pooled safety population, the most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were decreased white blood cell count (73%), nausea (72%), decreased hemoglobin (67%), decreased neutrophil count (65%), decreased lymphocyte count (60%), fatigue (55%), decreased platelet count (48%), increased aspartate

aminotransferase (46%), increased alanine aminotransferase (43%), increased blood alkaline phosphatase (39%), vomiting (38%), alopecia (37%), constipation (32%), decreased blood potassium (32%), decreased appetite (31%), diarrhea (30%), and musculoskeletal pain (24%).

ENHERTU in Combination with Pertuzumab

The pooled safety population reflects exposure to ENHERTU 5.4 mg/kg in combination with pertuzumab intravenously every 3 weeks in 431 patients in DESTINY-Breast07 (n=50), and DESTINY-Breast09 (n=381). Among these patients, 86% were exposed for >6 months and 73% were exposed for >1 year. In this pooled safety population, the most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were decreased white blood cell count (86%), decreased hemoglobin (80%), decreased neutrophil count (79%), nausea (74%), increased alanine aminotransferase (65%), diarrhea (64%), increased aspartate aminotransferase (63%), decreased lymphocyte count (61%), decreased platelet count (55%), increased blood alkaline phosphatase (54%), decreased blood potassium (54%), fatigue (53%), alopecia (48%), vomiting (46%), upper respiratory tract infection (32%), constipation (31%), decreased appetite (31%), decreased weight (28%), musculoskeletal pain (23%), increased blood bilirubin (23%) and abdominal pain (22%).

HER2-Positive Early Breast Cancer

DESTINY-Breast11

The safety of ENHERTU followed by THP was evaluated in 320 patients with HER2-positive (IHC 3+ or ISH+) early breast cancer who received at least 1 dose of ENHERTU 5.4 mg/kg followed by THP in DESTINY-Breast11. ENHERTU was administered by intravenous infusion once every three weeks for 4 cycles followed by THP for 4 cycles. The median duration of treatment was 5.6 months (range: 0.7 to 9.1) for patients who received ENHERTU followed by THP.

Serious adverse reactions occurred in 11% of patients receiving ENHERTU followed by THP, including COVID-19 (0.9%) and ILD/pneumonitis (0.6%). Fatal adverse reactions occurred in 0.6% of patients, including ILD/pneumonitis and death not otherwise specified (1 patient each).

In patients treated with ENHERTU followed by THP, the permanent discontinuation of ENHERTU due to adverse reactions occurred in 1.3%, of which ILD/pneumonitis accounted for 0.6%. Dose interruptions of ENHERTU due to adverse reactions occurred in 11% of patients. The most frequent adverse reactions ($>2\%$) associated with dose interruption were decreased neutrophil count and COVID-19. Dose reductions of ENHERTU occurred in 2.5% of patients treated with ENHERTU.

The most common ($\geq 20\%$) adverse reactions in patients treated with ENHERTU followed by THP, including laboratory abnormalities, were decreased hemoglobin (83%), increased alanine aminotransferase (79%), increased aspartate aminotransferase (74%), decreased white blood cell count (67%), nausea (65%), peripheral neuropathy (59%), diarrhea (59%), decreased neutrophil count (58%), alopecia (48%), fatigue (41%), decreased lymphocyte count (40%), rash (31%), musculoskeletal pain (30%), decreased blood potassium (29%), constipation (29%), vomiting (29%), stomatitis (23%), and decreased appetite (20%).

DESTINY-Breast05

The safety of ENHERTU was evaluated in 806 patients with HER2-positive breast cancer with residual invasive disease following neoadjuvant HER2-targeted therapy who then received at least one dose of ENHERTU 5.4 mg/kg. ENHERTU was administered by intravenous infusion once every three weeks for 14 cycles. The median duration of treatment was 10 months (range: 0.7 to 16) for patients who received ENHERTU.

Serious adverse reactions occurred in 17% of patients receiving ENHERTU. Serious adverse reactions in $\geq 1\%$ of patients who received ENHERTU were ILD/pneumonitis, radiation pneumonitis, pneumonia, and platelet count decreased. Fatal adverse reactions occurred in 0.4% of patients including ILD/pneumonitis (2 patients) and respiratory tract infection (1 patient).

Permanent discontinuation of ENHERTU due to an adverse reaction occurred in 18% of patients. The adverse reaction which resulted in permanent discontinuation of ENHERTU $>2\%$ included ILD/pneumonitis. Dose interruptions of ENHERTU due to an adverse reaction occurred in 50% of patients. Adverse reactions which required dosage interruptions in $>2\%$ included radiation pneumonitis, neutrophil count decreased, COVID-19, white blood cell count decreased, ILD/pneumonitis, platelet count decreased, upper respiratory tract infection, fatigue, cough, and pyrexia. Dose reductions of ENHERTU due to an adverse reaction occurred in 26% of

patients. Adverse reactions which required dose reductions in >2% of patients included nausea, fatigue, platelet count decreased, and neutrophil count decreased.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, in patients receiving ENHERTU were decreased white blood cell count (80%), decreased lymphocyte count (72%), decreased neutrophil count (72%), nausea (71%), decreased hemoglobin (61%), increased aspartate aminotransferase (60%), fatigue (54%), increased alanine aminotransferase (53%), decreased platelet count (46%), increased blood alkaline phosphatase (39%), constipation (32%), vomiting (31%), decreased blood potassium (27%), diarrhea (23%), musculoskeletal pain (23%), and decreased appetite (20%).

HER2-Positive Metastatic Breast Cancer

DESTINY-Breast09

The safety of ENHERTU 5.4 mg/kg in combination with pertuzumab was evaluated in DESTINY-Breast09, a randomized, three-arm, multicenter study including 763 patients with HER2-positive (IHC 3+ or ISH+) unresectable or metastatic breast cancer. Three hundred eighty-one patients received ENHERTU in combination with pertuzumab and 382 patients received THP (taxane [docetaxel or paclitaxel], trastuzumab, and pertuzumab). Among patients who received ENHERTU in combination with pertuzumab, the median duration of treatment was 22 months (range: 0.3 months to 44.5 months).

Serious adverse reactions occurred in 27% of patients receiving ENHERTU in combination with pertuzumab. Serious adverse reactions in >1% of patients were diarrhea, pneumonia, febrile neutropenia, hypokalemia, vomiting, ILD, pulmonary embolism, and sepsis. Fatalities due to adverse reactions occurred in 3.4% of patients including pneumonia (n=3), ILD (n=2), sepsis (n=2), pulmonary embolism, septic shock, acute kidney injury, dyspnea, febrile neutropenia, and intestinal ischemia (1 patient each).

ENHERTU was discontinued for adverse reactions in 21% of patients. The most frequent adverse reaction (>2%) associated with permanent discontinuation was ILD/pneumonitis (6%). Dose interruptions due to adverse reactions occurred in 69% of patients. The most frequent adverse reactions (>2%) associated with dose interruption were COVID-19, neutropenia, upper respiratory tract infection, fatigue, anemia, hypokalemia, ILD/pneumonitis, thrombocytopenia, pneumonia, diarrhea, transaminase increased, leukopenia, cough, pyrexia, decreased appetite, and blood bilirubin increased. Dose reductions occurred in 46% of patients treated with ENHERTU in combination with pertuzumab. The most frequent adverse reactions (>2%) associated with dose reduction were fatigue, neutropenia, nausea, diarrhea, ILD/pneumonitis, thrombocytopenia, vomiting, transaminases increased, decreased weight, febrile neutropenia, and hypokalemia.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were decreased white blood cell count (87%), decreased hemoglobin (80%), decreased neutrophil count (78%), nausea (75%), increased alanine aminotransferase (66%), diarrhea (64%), increased aspartate aminotransferase (62%), decreased lymphocyte count (62%), decreased platelet count (56%), increased blood alkaline phosphatase (55%), decreased blood potassium (54%), fatigue (53%), alopecia (48%), vomiting (46%), upper respiratory tract infection (33%), constipation (33%), decreased appetite (32%), decreased weight (30%), COVID-19 (28%), musculoskeletal pain (24%), increased blood bilirubin (23%), and abdominal pain (23%).

DESTINY-Breast03

The safety of ENHERTU was evaluated in 257 patients with unresectable or metastatic HER2-positive breast cancer who received at least 1 dose of ENHERTU 5.4 mg/kg intravenously once every 3 weeks in DESTINY-Breast03. The median duration of treatment was 14 months (range: 0.7 to 30) for patients who received ENHERTU.

Serious adverse reactions occurred in 19% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were vomiting, ILD, pneumonia, pyrexia, and urinary tract infection. Fatalities due to adverse reactions occurred in 0.8% of patients including COVID-19 and sudden death (1 patient each).

ENHERTU was permanently discontinued in 14% of patients, of which ILD/pneumonitis accounted for 8%. Dose interruptions due to adverse reactions occurred in 44% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose interruption were neutropenia, leukopenia, anemia, thrombocytopenia, pneumonia, nausea, fatigue, and ILD/pneumonitis. Dose reductions occurred in 21% of patients

treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were nausea, neutropenia, and fatigue.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were nausea (76%), decreased white blood cell count (74%), decreased neutrophil count (70%), increased aspartate aminotransferase (67%), decreased hemoglobin (64%), decreased lymphocyte count (55%), increased alanine aminotransferase (53%), decreased platelet count (52%), fatigue (49%), vomiting (49%), increased blood alkaline phosphatase (49%), alopecia (37%), decreased blood potassium (35%), constipation (34%), musculoskeletal pain (31%), diarrhea (29%), decreased appetite (29%), headache (22%), respiratory infection (22%), abdominal pain (21%), increased blood bilirubin (20%), and stomatitis (20%).

HER2-Low and HER2-Ultralow Metastatic Breast Cancer *DESTINY-Breast06*

The safety of ENHERTU was evaluated in 434 patients with unresectable or metastatic HER2-low (IHC 1+ or IHC 2+/ISH-) or HER2-ultralow (IHC 0 with membrane staining) breast cancer who received ENHERTU 5.4 mg/kg intravenously once every 3 weeks in DESTINY-Breast06. The median duration of treatment was 11 months (range: 0.4 to 39.6) for patients who received ENHERTU.

Serious adverse reactions occurred in 20% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were ILD/pneumonitis, COVID-19, febrile neutropenia, and hypokalemia. Fatalities due to adverse reactions occurred in 2.8% of patients including ILD (0.7%); sepsis (0.5%); and COVID-19 pneumonia, bacterial meningoencephalitis, neutropenic sepsis, peritonitis, cerebrovascular accident, general physical health deterioration (0.2% each).

ENHERTU was permanently discontinued in 14% of patients. The most frequent adverse reaction (>2%) associated with permanent discontinuation was ILD/pneumonitis. Dose interruptions due to adverse reactions occurred in 48% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose interruption were COVID-19, decreased neutrophil count, anemia, pyrexia, pneumonia, decreased white blood cell count, and ILD. Dose reductions occurred in 25% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were nausea, fatigue, decreased platelet count, and decreased neutrophil count.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were decreased white blood cell count (86%), decreased neutrophil count (75%), nausea (70%), decreased hemoglobin (69%), decreased lymphocyte count (66%), fatigue (53%), decreased platelet count (48%), alopecia (48%), increased alanine aminotransferase (44%), increased blood alkaline phosphatase (43%), increased aspartate aminotransferase (41%), decreased blood potassium (35%), diarrhea (34%), vomiting (34%), constipation (32%), decreased appetite (26%), COVID-19 (26%), and musculoskeletal pain (24%).

DESTINY-Breast04

The safety of ENHERTU was evaluated in 371 patients with unresectable or metastatic HER2-low (IHC 1+ or IHC 2+/ISH-) breast cancer who received ENHERTU 5.4 mg/kg intravenously once every 3 weeks in DESTINY-Breast04. The median duration of treatment was 8 months (range: 0.2 to 33) for patients who received ENHERTU.

Serious adverse reactions occurred in 28% of patients receiving ENHERTU. Serious adverse reactions in >1% of patients who received ENHERTU were ILD/pneumonitis, pneumonia, dyspnea, musculoskeletal pain, sepsis, anemia, febrile neutropenia, hypercalcemia, nausea, pyrexia, and vomiting. Fatalities due to adverse reactions occurred in 4% of patients including ILD/pneumonitis (3 patients); sepsis (2 patients); and ischemic colitis, disseminated intravascular coagulation, dyspnea, febrile neutropenia, general physical health deterioration, pleural effusion, and respiratory failure (1 patient each).

ENHERTU was permanently discontinued in 16% of patients, of which ILD/pneumonitis accounted for 8%. Dose interruptions due to adverse reactions occurred in 39% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose interruption were neutropenia, fatigue, anemia, leukopenia, COVID-19, ILD/pneumonitis, increased transaminases, and hyperbilirubinemia. Dose reductions occurred in 23% of patients treated with ENHERTU. The most frequent adverse reactions (>2%) associated with dose reduction were fatigue, nausea, thrombocytopenia, and neutropenia.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were nausea (76%), decreased white blood cell count (70%), decreased hemoglobin (64%), decreased neutrophil count (64%), decreased lymphocyte count (55%), fatigue (54%), decreased platelet count (44%), alopecia (40%), vomiting (40%), increased aspartate aminotransferase (38%), increased alanine aminotransferase (36%), constipation (34%), increased blood alkaline phosphatase (34%), decreased appetite (32%), musculoskeletal pain (32%), diarrhea (27%), and decreased blood potassium (25%).

HER2-Mutant Unresectable or Metastatic NSCLC (5.4 mg/kg)

DESTINY-Lung02 evaluated 2 dose levels (5.4 mg/kg [n=101] and 6.4 mg/kg [n=50]); however, only the results for the recommended dose of 5.4 mg/kg intravenously every 3 weeks are described below due to increased toxicity observed with the higher dose in patients with NSCLC, including ILD/pneumonitis.

The safety of ENHERTU was evaluated in 101 patients with HER2-mutant unresectable or metastatic NSCLC who received ENHERTU 5.4 mg/kg intravenously once every 3 weeks until disease progression or unacceptable toxicity in DESTINY-Lung02. The median duration of treatment was 8 months (range: 0.7 to 28) for patients who received ENHERTU.

Serious adverse reactions occurred in 40% of patients receiving ENHERTU. Serious adverse reactions in $>1\%$ of patients who received ENHERTU were ILD/pneumonitis, pleural effusion, thrombocytopenia, dyspnea, nausea, pneumonia, vomiting, myocarditis, pulmonary embolism, and increased troponin I. Fatalities due to adverse reactions occurred in 3% of patients including ILD/pneumonitis, cerebrovascular accident, and pneumococcal sepsis (1 patient each).

ENHERTU was permanently discontinued in 17% of patients. Adverse reactions which resulted in permanent discontinuation of ENHERTU were ILD/pneumonitis, pneumonia, blood bilirubin increased, hypokalemia, metastases to meninges, and myocarditis. Dose interruptions of ENHERTU due to adverse reactions occurred in 50% of patients. Adverse reactions which required dose interruption ($>2\%$) included neutropenia, COVID-19, ILD/pneumonitis, fatigue, anemia, and pneumonia. Dose reductions due to an adverse reaction occurred in 20% of patients. The most frequent adverse reactions ($>2\%$) associated with dose reduction were neutropenia, fatigue, and decreased appetite.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were decreased hemoglobin (68%), nausea (67%), decreased white blood cell count (66%), decreased neutrophil count (59%), decreased lymphocyte count (56%), increased aspartate aminotransferase (51%), decreased albumin (50%), decreased platelet count (49%), fatigue (48%), increased alanine aminotransferase (41%), decreased appetite (41%), constipation (38%), increased alkaline phosphatase (37%), vomiting (32%), decreased blood potassium (29%), diarrhea (24%), alopecia (22%), and musculoskeletal pain (21%).

HER2-Positive Locally Advanced or Metastatic Gastric Cancer (6.4 mg/kg)

The safety of ENHERTU was evaluated in 187 patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma in DESTINY-Gastric01. Patients intravenously received at least 1 dose of either ENHERTU (N=125) 6.4 mg/kg every 3 weeks or either irinotecan (N=55) 150 mg/m² biweekly or paclitaxel (N=7) 80 mg/m² weekly for 3 weeks. The median duration of treatment was 4.6 months (range: 0.7 to 22.3) for patients who received ENHERTU.

Serious adverse reactions occurred in 44% of patients receiving ENHERTU 6.4 mg/kg. Serious adverse reactions in $>2\%$ of patients who received ENHERTU were decreased appetite, ILD, anemia, dehydration, pneumonia, cholestatic jaundice, pyrexia, and tumor hemorrhage. Fatalities due to adverse reactions occurred in 2.4% of patients: disseminated intravascular coagulation, large intestine perforation, and pneumonia occurred in 1 patient each (0.8%).

ENHERTU was permanently discontinued in 15% of patients, of which ILD accounted for 6%. Dose interruptions due to adverse reactions occurred in 62% of patients treated with ENHERTU. The most frequent adverse reactions ($>2\%$) associated with dose interruption were neutropenia, anemia, decreased appetite, leukopenia, fatigue, thrombocytopenia, ILD, pneumonia, lymphopenia, upper respiratory tract infection, diarrhea, and decreased blood potassium. Dose reductions occurred in 32% of patients treated with ENHERTU. The most frequent adverse reactions ($>2\%$) associated with dose reduction were neutropenia, decreased appetite, fatigue, nausea, and febrile neutropenia.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were decreased hemoglobin (75%), decreased white blood cell count (74%), decreased neutrophil count (72%), decreased lymphocyte count (70%), decreased platelet count (68%), nausea (63%), decreased appetite (60%), increased aspartate aminotransferase (58%), fatigue (55%), increased blood alkaline phosphatase (54%), increased alanine aminotransferase (47%), diarrhea (32%), decreased blood potassium (30%), vomiting (26%), constipation (24%), increased blood bilirubin (24%), pyrexia (24%), and alopecia (22%).

HER2-Positive (IHC 3+) Unresectable or Metastatic Solid Tumors

The safety of ENHERTU was evaluated in 347 adult patients with unresectable or metastatic HER2-positive (IHC 3+) solid tumors who received ENHERTU 5.4 mg/kg intravenously once every 3 weeks in DESTINY-Breast01, DESTINY-PanTumor02, DESTINY-Lung01, and DESTINY-CRC02. The median duration of treatment was 8.3 months (range 0.7 to 30.2).

Serious adverse reactions occurred in 34% of patients receiving ENHERTU. Serious adverse reactions in $>1\%$ of patients who received ENHERTU were sepsis, pneumonia, vomiting, urinary tract infection, abdominal pain, nausea, pneumonitis, pleural effusion, hemorrhage, COVID-19, fatigue, acute kidney injury, anemia, cellulitis, and dyspnea. Fatalities due to adverse reactions occurred in 6.3% of patients including ILD/pneumonitis (2.3%), cardiac arrest (0.6%), COVID-19 (0.6%), and sepsis (0.6%). The following events occurred in 1 patient each (0.3%): acute kidney injury, cerebrovascular accident, general physical health deterioration, pneumonia, and hemorrhagic shock.

ENHERTU was permanently discontinued in 15% of patients, of which ILD/pneumonitis accounted for 10%. Dose interruptions due to adverse reactions occurred in 48% of patients. The most frequent adverse reactions ($>2\%$) associated with dose interruption were decreased neutrophil count, anemia, COVID-19, fatigue, decreased white blood cell count, and ILD/pneumonitis. Dose reductions occurred in 27% of patients treated with ENHERTU. The most frequent adverse reactions ($>2\%$) associated with dose reduction were fatigue, nausea, decreased neutrophil count, ILD/pneumonitis, and diarrhea.

The most common ($\geq 20\%$) adverse reactions, including laboratory abnormalities, were decreased white blood cell count (75%), nausea (69%), decreased hemoglobin (67%), decreased neutrophil count (66%), fatigue (59%), decreased lymphocyte count (58%), decreased platelet count (51%), increased aspartate aminotransferase (45%), increased alanine aminotransferase (44%), increased blood alkaline phosphatase (36%), vomiting (35%), decreased appetite (34%), alopecia (34%), diarrhea (31%), decreased blood potassium (29%), constipation (28%), decreased sodium (22%), stomatitis (20%), and upper respiratory tract infection (20%).

Use in Specific Populations

- **Pregnancy:** ENHERTU can cause fetal harm when administered to a pregnant woman. Advise patients of the potential risks to a fetus. There are clinical considerations if ENHERTU is used in pregnant women, or if a patient becomes pregnant within 7 months after the last dose of ENHERTU.
- **Lactation:** There are no data regarding the presence of ENHERTU in human milk, the effects on the breastfed child, or the effects on milk production. Because of the potential for serious adverse reactions in a breastfed child, advise women not to breastfeed during treatment with ENHERTU and for 7 months after the last dose.
- **Females and Males of Reproductive Potential:** Pregnancy testing: Verify pregnancy status of females of reproductive potential prior to initiation of ENHERTU. Contraception: *Females:* ENHERTU can cause fetal harm when administered to a pregnant woman. Advise females of reproductive potential to use effective contraception during treatment with ENHERTU and for 7 months after the last dose. *Males:* Advise male patients with female partners of reproductive potential to use effective contraception during treatment with ENHERTU and for 4 months after the last dose. Infertility: ENHERTU may impair male reproductive function and fertility.
- **Pediatric Use:** Safety and effectiveness of ENHERTU have not been established in pediatric patients.
- **Geriatric Use:** *ENHERTU as Monotherapy:* Of the 2233 patients treated with ENHERTU 5.4 mg/kg, 28% were ≥ 65 years and 6% were ≥ 75 years. No overall differences in efficacy within clinical studies were observed between patients ≥ 65 years compared to younger patients. There was a higher incidence of Grade 3-4 adverse reactions observed in patients aged ≥ 65 years (56%) as compared to younger patients (49%). Of the 125 patients with HER2-positive locally advanced or metastatic gastric or GEJ adenocarcinoma treated with ENHERTU 6.4 mg/kg in DESTINY-Gastric01, 56% were ≥ 65 years and 14% were ≥ 75 years. No overall

differences in efficacy or safety were observed between patients ≥ 65 years of age compared to younger patients. *ENHERTU in Combination with Pertuzumab*: In patients with HER2-positive unresectable or metastatic breast cancer treated with ENHERTU 5.4 mg/kg in combination with pertuzumab (N=431), 17% were ≥ 65 years and 3% were ≥ 75 years. No overall differences in efficacy or safety were observed between patients ≥ 65 years compared to younger patients. *ENHERTU followed by THP*: Of the 320 patients with HER2-positive early breast cancer treated with ENHERTU 5.4 mg/kg followed by THP, 12% were ≥ 65 years and 1.6% were ≥ 75 years. No overall differences in efficacy were observed between patients ≥ 65 years compared to younger patients. There was a higher incidence of Grade 3-4 adverse reactions observed in patients ≥ 65 years (38%) as compared to younger patients (30%).

- **Renal Impairment:** A higher incidence of Grade 1 and 2 ILD/pneumonitis has been observed in patients with moderate renal impairment. Monitor patients with moderate renal impairment more frequently. The recommended dosage of ENHERTU has not been established for patients with severe renal impairment (CLCr < 30 mL/min).
- **Hepatic Impairment:** In patients with moderate hepatic impairment, due to potentially increased exposure, monitor for increased adverse reactions related to the topoisomerase inhibitor, DXd. The recommended dosage of ENHERTU has not been established for patients with severe hepatic impairment (total bilirubin > 3 times ULN and any AST).

To report SUSPECTED ADVERSE REACTIONS, contact Daiichi Sankyo, Inc. at 1-877-437-7763 or FDA at 1-800-FDA-1088 or [fda.gov/medwatch](https://www.fda.gov/medwatch).

Please see accompanying full [Prescribing Information](#), including **Boxed WARNINGS, and [Medication Guide](#).**

About the ADC Portfolio of Daiichi Sankyo

The Daiichi Sankyo ADC portfolio consists of nine ADCs in clinical development crafted from ADC technology discovered in-house by Daiichi Sankyo.

The DXd ADC Technology platform of Daiichi Sankyo consists of seven ADCs in clinical development where each ADC is comprised of a monoclonal antibody attached to a number of topoisomerase I inhibitor payloads (an exatecan derivative, DXd) via tetrapeptide-based cleavable linkers. The DXd ADCs include Enhertu and Datroway, which are being jointly developed and commercialized globally with AstraZeneca, and ifinatamab deruxtecan (I-DXd), raludotatug deruxtecan (R-DXd) and patritumab deruxtecan (HER3-DXd), which are being jointly developed and commercialized globally with Merck & Co., Inc., Rahway, NJ, USA. DS-3939 and DS3790 are being developed by Daiichi Sankyo.

Additional ADCs being developed by Daiichi Sankyo include DS3610, which consists of an antibody attached to a novel payload that acts as an agonist of STING, and DS1025, which consists of a CD25 directed antibody attached to an immuno-oncology optimized cytotoxic payload.

Ifinatamab deruxtecan, raludotatug deruxtecan, patritumab deruxtecan, DS-3939, DS3610, DS3790 and DS1025 are investigational medicines that have not been approved for any indication in any country. Safety and efficacy have not been established.

About Daiichi Sankyo

Daiichi Sankyo (TSE: 4568) is a global healthcare company committed to being a trusted healthcare innovator, contributing to the enrichment of quality of life around the world through its strengths in science and technology. At Daiichi Sankyo, we believe progress depends on asking more of science, challenging assumptions and pursuing better answers. Progress like this is built by people who never stop learning, questioning and pushing what's possible. Together, we work to bring new breakthrough medicines within reach for patients and help shape what comes next in oncology and beyond. For more information, please visit www.daiichisankyo.com and connect with us on [LinkedIn](#).

MEDIA CONTACTS:**Global:**

Jennifer Brennan

[*jennifer.brennan@daiichisankyo.com*](mailto:jennifer.brennan@daiichisankyo.com)

+1 908 900 3183 (mobile)

Japan:

[*DS-PR_jp@daiichisankyo.com*](mailto:DS-PR_jp@daiichisankyo.com)

INVESTOR RELATIONS CONTACT:

[*DaiichiSankyoIR_jp@daiichisankyo.com*](mailto:DaiichiSankyoIR_jp@daiichisankyo.com)